

Health Newsletter

-Generali China GBD

2026 September



往期中意健康月刊

Contents

P3

September Health Begins
by Reading the Signals

P4

World Alzheimer's Day: Protecting
Brian Starts in Midlife

P8

Blood Lipid Puzzles and
Help You Better Manage Your Blood
Lipid

P11

What Causes
a Heart Attack

P15

Six Practical Moves for Stronger
Bones

P18

Why Are You Drawn to Brokenness
The Psychology of the Rescuer Role



September Health Begins by Reading the Signals

September offers a natural opportunity to reset our routines and listen more closely to what our bodies are telling us. Health is more than the absence of discomfort, and it does not begin only after a problem appears. It is an ongoing practice of noticing change, understanding risk, and taking appropriate action at the right time.

The body and mind rarely remain completely silent. Changes in memory and cognition, health-check results and heart rhythm, bone strength and mobility, or a recurring sense of depletion in a relationship can all send different kinds of signals. Some deserve closer attention, while others point to habits that can be adjusted early. This issue connects these seemingly separate clues and explores the longer-term patterns behind them.

Being proactive does not mean becoming anxious about every number or changing everything at once. It may begin with one honest observation: understand what a test result means, ask a qualified professional a question, make room for movement and sleep, or set a clearer emotional boundary. Sustainable health management is usually built through small, practical choices that can be repeated.

This September, may we meet change with greater understanding and less neglect, then turn what we notice into action that fits our own lives. GCL is here to support the long-term wellbeing of your body, your heart and brain, and your inner life.

[Stay alert to cardiovascular and cerebrovascular disease risks and learn more about targeted coverage from 中意意安护 →](#)



Scan for more info



World Alzheimer's Day: Protecting Brian Starts in Midlife



Every year on September 21, we observe World Alzheimer's Day. Alzheimer's disease is commonly referred to as "senile dementia" or simply "dementia." When people hear "senile dementia," many assume it's a concern only for 他 the elderly. **However, scientific research shows that the battle to protect our brains begins in midlife, or even earlier.**



“Brain Crisis” Starts from Midlife.

We commonly think of dementia as an “old age disease” and often overlook its long incubation period. In fact, **the pathological changes of Alzheimer’s disease may begin 20 to 30 years before symptoms appear.** This means that while you are still building your career and working late nights, your brain may already be undergoing subtle changes.

Unhealthy lifestyles in midlife are significant drivers of cognitive decline. Chronic sleep deprivation, a sedentary lifestyle, high-fat and high-sugar diets, social isolation, and the prevalence of “three highs” (hypertension, hyperglycemia, and hyperlipidemia) have greatly increased the risk of cognitive decline among middle-aged populations. **Studies confirm that scientific prevention can reduce the risk of dementia by up to 45%, and the critical window for intervention is in midlife.**

Common Cognitive Misconceptions Ignoring Controllable Factors:

Beyond unchangeable age and genetics, **more than one-third of dementia cases are linked to modifiable lifestyle factors.**

Poor management of chronic diseases, unhealthy diets, lack of exercise, and smoking are all hidden “brain killers.”

Blindly Choosing Supplements: The market is flooded with various brain supplements. Research shows that while **specific nutritional supplements (such as polyphenols and probiotics)** may offer modest cognitive benefits in the early stages of the disease, their effects are often limited, and **they should not replace a healthy lifestyle.**

Ignoring Early Warning Signs :

Many people dismiss memory decline as a normal part of aging and miss the optimal window for intervention. **The early diagnosis rate is less than 20%,** and most patients are already in the moderate to severe stages when diagnosed.

Start Scientific Intervention Now

Prevention is better than cure. Starting in midlife, we should build a brain shield from the following aspects:

Manage the Chronic

Diseases: Abnormalities in blood pressure, blood sugar, and blood lipids directly damage cerebral blood vessels. Regular monitoring and taking medication as prescribed by a doctor are the cornerstones of preventing vascular dementia and Alzheimer's disease.

Adopt the “MIND” Brain-Healthy

Diet: The “MIND diet,” which combines the Mediterranean and DASH diets, has been proven to significantly slow brain aging. The core principle is: Eat more green leafy vegetables, berries (blueberries, strawberries), nuts, whole grains, fish, and poultry; eat less red meat, butter, fried foods, and sweets.

Get Moving for a More Active Brain :

Regular aerobic exercise (e.g., brisk walking, jogging, swimming) promotes cerebral blood flow and neurogenesis. At least 150 minutes of moderate-intensity exercise per week, combined with two sessions of strength training, keep your brain more active.

Keep Active Socializing and Learning :

Persist in reading, learning new skills, and engaging in puzzle games to build cognitive reserve. At the same time, communicate more with others and participate in community activities to avoid prolonged isolation. Social isolation has been listed by The Lancet as a major risk factor for dementia.





About Supplements: To Take or Not to Take?

First, supplements should never replace basic lifestyle interventions like the MIND diet and exercise. If you have a balanced diet and normal physical indicators, there is no need for extra supplementation. If you do have a need, **you should do it under the guidance of a doctor or nutritionist with following principles.**

1. Check nutritional deficiencies first, avoid blind supplementation

Prioritize blood tests for vitamin D, vitamin B12, folate and homocysteine. Supplements should be administered under the guidance of a nutritionist when a deficiency is confirmed. Do not take multiple brainhealth complex products simultaneously.

2. Foodfirst approach, supplements used as an aid

Obtain Omega3 (DHA), polyphenols and probiotics from daily meals whenever possible. Get DHA from fatty fish and nuts, polyphenols from berries and darkcolored vegetables, and probiotics from unsweetened fermented foods. Consider supplements when diet cannot meet intake requirements.

3. Scientifically regulate the gutbrain axis

Gut health is linked to brain function. Studies show that certain phytochemicals (e.g., polyphenols) and probiotics may exert beneficial effects via the gutbrain axis. For probiotics, choose strains of Bifidobacterium and Lactobacillus. When supplementing with polyphenols such as curcumin, opt for easilyabsorbed formulations (e.g., liposomal forms). Take them with oilcontaining main meals to boost absorption and avoid over-dose intake.



Early Recognition, Seek Medical Attention Promptly

It is recommended to visit a neurology clinic when the following situations occur:

- **You just put something down and immediately forget where it is**, and this happens with increasing frequency.
- **You feel unfamiliar with familiar environments or tasks**, such as getting lost on a route you know well or being unable to use familiar appliances.
- **Changes in personality and behavior**, becoming suspicious, anxious, irritable, or apathetic.

Blood Lipid Puzzles and Help You Better Manage Your Blood Lipid



The prevalence of dyslipidemia among Chinese adults has nearly reached two fifth, meaning one in three adults has unhealthy lipid levels. Hyperlipidemia is a direct culprit behind heart attacks and strokes, yet the public holds many misconceptions—such as believing that simply avoiding fatty meat is enough. Today, we’ ll guide you out of these myths and help you manage your blood lipids scientifically.

The Four Lipid Indicators: Reading Your Report Easily

The “four lipid indicators” on your health check report can be utterly confusing, and it is not so hard to interpret the report. When you get your lipid panel results, focus on these four key indicators:

Total Cholesterol (TC): The sum of all cholesterol in your blood **and it provides the raw material for atherosclerosis.** The ideal value is below 5.2 mmol/L.

Triglycerides (TG): Primarily reflects fat reserves and energy metabolism. **It is most closely related to diet;** overeating and heavy drinking can cause it to spike. The ideal value is below 1.7 mmol/L.

Low-Density Lipoprotein Cholesterol (LDL-C): **This is what we commonly call the “bad cholesterol.”** It is the main culprit behind clogged arteries and plaque formation. For high-risk individuals, **LDL-C is the primary target for control.** The ideal value is generally below 3.4 mmol/L, but patients with established cardiovascular disease need much lower levels (<1.8 mmol/L or even <1.4 mmol/L).

High-Density Lipoprotein Cholesterol (HDL-C): **This is the “good cholesterol.”** It collects excess cholesterol from peripheral tissues of the body and transports it back to the liver for metabolism and excretion, **helping protect blood vessels.** **The higher the value, the better;** ideal value ≥ 1.04 mmol/L.





Common Misconceptions on Blood Lipid

Misconception 1: High cholesterol only affects overweight people; thin people don't need to worry.

This is not true. Thin individuals can also have dyslipidemia due to genetics, high metabolic stress, or hidden visceral fat. Therefore, regardless of body type, **adults should have their lipid levels checked at least once a year.**

Misconception 2: If you don't eat meat and only eat vegetarian food, your blood lipids will be normal.

This is the most common myth. **Refined carbohydrates (rice, noodles, steamed buns, bread) and sugary drinks are converted into triglycerides in the body, also raising lipid levels.** Furthermore, if vegetarians use too much cooking oil or consume large amounts of vegetable oils, excess calories can still elevate lipids.

Misconception 3: All vegetable oils are healthy and can be consumed freely.

Not all vegetable oils are good. Coconut oil and palm oil are rich in saturated fatty acids, similar to animal fats. Moreover, **vegetable oils that are repeatedly fried or heated at high temperatures produce trans fatty acids, which will accelerate atherosclerosis.** It is recommended to choose **oils rich in unsaturated fatty acids, such as olive oil, camellia oil, and flaxseed oil, and limit daily intake to 25–30 grams.**

Misconception 4: High blood lipids have no symptoms, so there's no need to take action.

In its early stages, hyperlipidemia indeed has no obvious symptoms, but it is silently damaging the vascular endothelium and accelerating atherosclerosis. **When a heart attack or stroke occurs, it is already too late.**



Lipid-Lowering Medications: When to Start and How to Take Them?

Many people agonize over whether to take medication, fearing that “once you start, you can never stop.” This fear is unnecessary.

When is medication therapy needed?

For patients **already diagnosed with cardiovascular disease (coronary heart disease, stroke) or diabetes with high-risk factors**, regardless of their current lipid levels, statin therapy should be initiated, and **LDL-C needs to be lowered to at least 1.8 mmol/L**.

For those **without a history of cardiovascular disease**, if **LDL-C remains ≥ 3.4 mmol/L after 3 to 6 months of lifestyle intervention**, or if **other risk factors (hypertension, smoking, obesity, etc.) are present**, medication therapy should also be considered.

Common Treatment Plans and Precautions:

Statin is the common medicine of lipid-lowering therapy and significantly reduce the risk of cardiovascular events.


Medication is **usually taken after dinner or at bedtime**, as cholesterol synthesis is most active at night.

Regular follow-up testing of lipids, liver function (ALT), and muscle-related markers (CK) is required during treatment. If muscle aches, fatigue, or dark urine occur, seek medical attention promptly.

Statins require long-term adherence; stopping them abruptly can cause lipid levels to rebound and increase risk.

What Causes a Heart Attack

September 29 marks World Heart Day, highlighting the importance of heart health education for all. If you're concerned about a heart attack, the best thing you can do is understand the causes and risk factors, so you can take measures to lower your chances of having one. Heart attacks are life-threatening and can lead to heart failure. While atherosclerosis (buildup of plaque in the arteries) is the most common cause of a heart attack, several other lifestyle habits and underlying conditions, including having a family history of heart attacks, can also raise your risk.





Risk Factors

Researchers have identified many risk factors for heart disease. Some are modifiable, meaning you have the power to change them, whereas others, like age and genetics, are considered non-modifiable.

Age: Heart attacks can happen to anyone of any age or sex but are more likely as you get older, particularly in males over 45 and females over 55.

Atherosclerosis: Atherosclerosis involves the buildup of cholesterol plaques in your blood vessels. If these plaques become disrupted, they can trigger the body's blood clotting mechanism, leading to a blood clot. This clot can block blood flow in a coronary artery, potentially causing a heart attack.

Diabetes: According to the National Institute of Diabetes and Digestive and Kidney Disorders, adults with diabetes are about twice as likely as those without diabetes to experience a heart attack or stroke at some point in their life.

Abnormal Cholesterol: Abnormal cholesterol, or dyslipidemia, is another major risk factor for heart disease that also may not cause symptoms. High levels of low-density lipoprotein (LDL) cholesterol, also known as "bad cholesterol", can increase heart attack risk.

High Blood Pressure : High blood pressure (hypertension) is elevated pressure in your arteries. The condition is strongly linked to cardiovascular diseases, including heart attacks and strokes. It's important to see your healthcare provider for high blood pressure screening since it normally doesn't cause symptoms, and you may not otherwise know you have it.

Lifestyle Habits: Your lifestyle habits greatly influence your risk of future heart attacks. You may have a higher risk of developing a heart attack if you follow any of these lifestyle habits:

- Having a sedentary lifestyle without regular physical activity or exercise
- Smoking cigarettes
- Eating a diet high in ultra-processed foods, salt, trans fats, and saturated fats
- Getting poor quality or not enough sleep
- Not managing stress levels appropriately

High Blood Pressure : High blood pressure (hypertension) is elevated pressure in your arteries. The condition is strongly linked to cardiovascular diseases, including heart attacks and strokes. It's important to see your healthcare provider for high blood pressure screening since it normally doesn't cause symptoms, and you may not otherwise know you have it.

Lifestyle Habits: Your lifestyle habits greatly influence your risk of future heart attacks. You may have a higher risk of developing a heart attack if you follow any of these lifestyle habits:

- Having a sedentary lifestyle without regular physical activity or exercise
- Smoking cigarettes
- Eating a diet high in ultra-processed foods, salt, trans fats, and saturated fats
- Getting poor quality or not enough sleep
- Not managing stress levels appropriately



Daily Prevention & Wellness

Avoid Common Triggers: Sleep deprivation, excessive strong tea/coffee, heavy alcohol, and drastic emotional fluctuations are common triggers.

Recommendations: caffeine ≤ 400 mg/day (2 cups of Americano), 7–8 hours sleep per day, limit alcohol, practice stress management.

Manage Chronic Diseases: Hypertension, diabetes, and dyslipidemia are independent risk factors for common arrhythmias, and sudden cardiac death. High-risk individuals should monitor these parameters regularly, adhere to medications, and **never discontinue antihypertensives, glucose-lowering drugs, or statins without guidance.**

Balanced Diet & Aerobic Exercise: Adopt a **Mediterranean diet** (vegetables, fruits, whole grains, fish, olive oil), limit saturated fats. Engage in **moderate-intensity aerobic exercise** (brisk walking,

swimming, cycling) for **≥ 150 min/week** to improve autonomic function. Those with structural heart disease should exercise under supervision and **avoid high-intensity or competitive sports.**

Regular Follow-up & Self-Monitoring: Patients with confirmed arrhythmias should have **cardiology follow-up every 3–6 months.** Smart wearables (e.g., smart watches) can assist home monitoring, but **device alerts cannot replace professional diagnosis.** All abnormal signals require clinical confirmation.





Six Practical Moves for Stronger Bones

Bone loss often develops without obvious symptoms. Its impact may first become apparent after a fall and fracture, a decline in mobility, or a growing fear of being active. Bone health is not determined by bone density alone. Muscle strength, balance, reaction time, and the safety of the home environment also matter. Regular weight-bearing, resistance, and balance exercise can address several of these factors together. The following six movements require little equipment and can provide a practical foundation for home exercise.

The goal is not to do as much as possible in a single session. It is to give the body an appropriate stimulus repeatedly and within a manageable range. Steady technique, long-term consistency, and gradual progression usually matter more than speed, a large range of motion, or heavy resistance.

Requires final confirmation by you or a medical colleague: People with diagnosed osteoporosis, a previous fragility fracture, a recent fall, significant pain, dizziness, or balance problems should be assessed by a physician or physical therapist before adjusting the exercises, resistance, or training volume.

Why Bones Need the Stimulus of Movement

Bone is not a fixed framework. It is living tissue that is continuously renewed. Weight bearing, muscle contraction, and appropriate external loads provide mechanical stimulation to the skeleton, while the muscles and nervous system learn to control the body more steadily. With age, less daily movement, prolonged sitting, or avoiding activity because of pain or fear of falling can reduce this stimulus and contribute to further losses in strength and balance.

A bone-supporting exercise plan should therefore address at least three areas. Weight-bearing activities, such as walking, stair climbing, and dancing, make the body work against gravity while upright. Each has a different role, and the combination more closely reflects the demands of daily life.

Why Strength and Balance Training Belong Together

Fracture risk is not determined by bone mass alone. Limited lower-body strength can make it harder to rise from a chair, negotiate steps, or move around an obstacle. Poor ankle control and balance can make a loss of footing more likely. Walking helps maintain activity, but it may not provide enough resistance for every major muscle group. A plan that includes strength, weight-bearing activity, and balance better supports the abilities needed for independent living.

For many people, a suitable starting point is one in which every repetition can be controlled, breathing remains normal, and discomfort does not continue to worsen the next day. Once a movement feels familiar, progress by changing one factor at a time: add repetitions, slow the lowering phase, use slightly less hand support, or add light resistance. Do not change everything at once.



Three Checks Before You Begin

First, check the environment. Use a sturdy chair without wheels and place it against a wall or where it cannot slide. Wear well-fitting shoes with nonslip soles, and remove cords, loose rug edges, and clutter from the floor. If an exercise requires a wall, rail, or countertop, make sure the support will not move.

Next, check how you feel that day. Pause exercise if you have a fever, unusual fatigue, chest discomfort, dizziness, or an acute flare of joint or back pain. Warm up with three to five minutes of easy walking or marching in place, followed by gentle movement of the shoulders, hips, and ankles. Finally, check the quality of each repetition. Muscles may feel as though they are working, but there should be no sharp pain, numbness, or sense that you are about to fall. Keep breathing normally and avoid holding your breath.





Supported Squat

This movement primarily trains the hips and thighs. It also resembles everyday tasks such as sitting down, rising from a chair, and reaching toward a lower surface.

Stand with your feet about hip-width apart and hold a stable countertop or chair back. Send your hips backward, then bend your knees as if you were sitting down. Keep your chest comfortably lifted and your knees traveling in roughly the same direction as your toes. Press through your feet to stand. Only lower as far as your knees, hips, and lower back feel comfortable.

When beginning, place a chair behind you and lightly touch the seat before standing. Common problems include letting the knees collapse inward, dropping suddenly, or bending far forward to reach a lower position. Begin with a small range and increase the depth or reduce hand support only after you can repeat the movement steadily.

Standing Heel Raise

Heel raises train the calf muscles and ankle control. These abilities help propel the body during walking, support stair climbing, and assist with recovering from small shifts in balance.

Stand with your feet parallel and hold a stable support. Slowly lift your heels so your weight moves toward the balls of your feet while the toes remain relaxed. Pause briefly, then lower your heels with control. Do not bounce, and do not allow the ankles to roll markedly inward or outward.

Begin with both feet working together. If the movement is steady, lengthen the lowering phase or gradually rely less on your hands. A single-leg heel raise is much more demanding and should not be attempted simply because the two-leg version feels easy, particularly if you have balance problems or a recent fall.



Low Step-Up

A step-up combines single-leg strength, weight transfer, and balance. It is similar to the demands of stairs, curbs, and thresholds.

Use a low, stable step near a handrail or wall. Place one whole foot on the step, shift your weight forward and upward, and bring the other foot up. Return to the floor slowly, one foot at a time. Alternate the leading leg, place each foot securely, and prioritize control over speed.

If the knee is uncomfortable, use a lower step or begin by practicing forward and backward weight shifts on level ground. Do not use a loose footstool or place the step on a rug. Progress by improving control or adding repetitions before choosing a higher step.

Resistance Band Row

Rows train the upper back, the muscles around the shoulder blades, and the arms. These areas contribute to carrying objects, opening doors, and maintaining the upper-body posture used in daily tasks.

Sit or stand tall. Secure the resistance band to a reliable anchor or use it safely according to the product instructions. Hold the band with your wrists in a neutral position, draw your elbows backward, and gently bring your shoulder blades toward each other. Return slowly. Keep your neck relaxed and avoid shrugging or leaning back.

Begin with a light band that has no cracks or signs of deterioration, and stay out of the path the band would travel if it slipped. There is no need to pinch the shoulder blades forcefully at the end of the movement. Once the trunk remains steady, increase either the resistance or the repetitions by a small amount.



Supported Side Leg Raise

A side leg raise primarily trains the muscles along the outside of the hip. Hip control supports single-leg stance, sideways steps, and pelvic stability while walking.

Stand upright while holding a countertop or chair back. Lift one leg a short distance to the side with the toes pointing forward, pause, then lower it slowly. Do not lock the knee of the standing leg. Keep the trunk upright and avoid turning the toes to create the movement. Train both sides.


The leg does not need to lift high; feeling the outside of the hip working is enough. A noticeable sideways lean usually means that the range is too large or control is limited. With practice, reduce hand support or add light resistance only if a qualified professional has advised it.

Wall Push-Up

A wall push-up trains the chest, shoulders, and arms. It is easier to control than a floor push-up and can be adjusted to suit different ability levels.

Stand facing a wall and place your hands near shoulder height, slightly wider than the shoulders. Step back a short distance. Bend your elbows to bring your chest toward the wall, then push back to the starting position. Keep a comfortable straight line from the head through the legs, gently engage the abdomen, and avoid letting the lower back sag or the head jut forward.

Standing farther from the wall usually makes the movement harder. Choose a position that allows you to remain stable throughout; your chest does not have to touch the wall. If the shoulder hurts, adjust the hand position and elbow angle. Stop and seek an assessment if discomfort continues.



Fitting the Six Moves Into Your Week

For the first session, try five to eight repetitions of each movement, rest briefly between exercises, and complete one round. If the next day brings only mild muscle soreness and ordinary activities are unaffected, maintain the same amount or increase it slightly at the following session. If pain persists, worsens, or interferes with walking or sleep, reduce the training and identify the cause. Repetition counts are only a guide; movement quality and the body's response matter more.

Once the movements are familiar, divide them across two or three strength sessions each week and allow recovery time between sessions. Older adults should also include activities combining balance and strength on three or more days per week. Balance practice can be simple: stand heel to toe beside a stable support, walk slowly along a straight path, or practice controlled changes of direction. Anyone at clear risk of falling should not independently attempt difficult drills such as standing with the eyes closed or removing all support.

Brisk walking, stair climbing, and dancing can add regular weight-bearing activity and support cardiovascular fitness. Swimming and cycling may be easier on the joints and can improve fitness, but they do not load the body in the same way as walking. They should not be treated as the entire bone-strengthening program. Combining activities according to joint comfort and personal interest can make the plan easier to maintain.

Why Are You Drawn to Brokenness

The Psychology of the Rescuer Role

In the film *Memories of Matsuko*, Matsuko grows up trying to make her father smile and later enters one painful relationship after another. She is repeatedly drawn to people who seem lonely, wounded, or in need of rescue. She tries to secure love by giving more, fixing more, and asking for less. The story is extreme, but the feeling behind it is familiar to many people: you know a relationship is wearing you down, yet you still feel compelled to care for the other person, repair the damage, or treat leaving as a form of betrayal.

People sometimes describe this as an attraction to “brokenness.” The word broken is not a diagnosis and should not be used to reduce a person to a label. Here, it refers to vulnerability, trauma, loss, or instability that we perceive in someone else. Being moved by vulnerability is not itself unhealthy. The concern begins when empathy turns into obligation, love must be proven through sacrifice, and your boundaries, safety, and needs remain last on the list.





Empathy Draws Us Toward Another Person's Pain

Human beings are equipped to notice and respond to other people's emotions. Affection Exchange Theory describes giving and receiving affection as important parts of human connection. Research on empathy also suggests that seeing another person in pain recruits brain networks involved in emotion, bodily awareness, and preparation for action. We do not literally reproduce the other person's entire experience, but we may have a partial, embodied response that helps us understand what they are going through.

Our own history shapes that response. If you have lived through similar rejection, neglect, or insecurity, another person's story may feel especially compelling. The Perception-Action Model proposes that we use our own distributed representations and past experience to understand another person's state. Familiarity, similarity, and prior experience can all intensify empathic responding. The other person's pain may then feel less like information and more like a demand: I need to do something.

There is nothing shameful about that impulse. Listening, staying present, and offering support within your capacity are healthy parts of a relationship. But empathy does not automatically make you responsible for repairing another adult, nor does it prove that the two of you are compatible. Understanding someone and assuming responsibility for that person's choices and emotions are different things.

Compulsive Caregiving Can Repeat a Familiar Script

Why do some people consistently become the caregiver while finding it difficult to receive care? Attachment literature has used the term compulsive caregiving for an imbalanced pattern in which a person repeatedly comforts, provides, and solves problems while suppressing their own need for support. Giving may feel like the safest, or only familiar, way to remain close to another person.

This pattern may be related to role reversal in childhood. A child who should be receiving reliable care may instead have to manage a parent's illness, emotional fragility, addiction, or absence. Over time, the child may absorb unspoken rules: people stay only when I am useful; my needs will burden others; if I work harder, I can control what happens to the relationship.

Not every generous caregiver has a history of childhood trauma, and not everyone who experienced role reversal repeats the same relationship pattern. Psychological concepts are clues for reflection, not verdicts about an individual. More useful questions are whether giving is freely chosen, whether care can flow in both directions, and whether saying no or taking a break triggers intense guilt or fear.

When self-worth depends on being needed, a vulnerable and unstable partner can feel unusually compelling. Rescue offers an apparent chance to rewrite an old script: This time, if I do enough, the ending will be different. Yet another adult's willingness to seek help, change behavior, and accept responsibility cannot be determined by how much the caregiver sacrifices.



Why Unpredictable Rewards Make It Harder to Leave

Behavioral research has shown that when reinforcement arrives intermittently and unpredictably, an established behavior can become especially resistant to stopping. Experimental psychology calls one form of this arrangement variable-ratio reinforcement. The concept is often illustrated with gambling, but it can also help explain a relationship experience: long stretches of distance, rejection, or broken promises are interrupted by a rare moment of tenderness, apology, or dependence, which renews the desire to keep trying.

That unusual moment of closeness may be interpreted as the return of the person's "real self," while earlier harm is treated as temporary. Hoping to receive the same response again, a person may give more, expect less, and rationalize further discomfort. This should not be reduced to a simple claim that one neurotransmitter is constantly activated. The more important point is that uncertainty, hope, and previous investment can combine to make disengagement difficult.

Caring for someone can also create an immediate sense of control and worth. When the rest of life feels uncertain, organizing another person's life, managing consequences, or responding to repeated crises can make the belief "I am useful" or "I can change the outcome" feel concrete. Helping is not the problem. The problem is relying on being needed as the only source of value, or repeatedly shielding the other person from responsibilities that belong to them.

Look at the overall pattern rather than a few high points. Are promises kept consistently? Is there genuine repair after conflict? Does the other person respect your no? Can you admit that you are tired without being punished, ignored, or threatened? The quality of a relationship is best judged by repeated interactions over time, not by asking a handful of sweet moments to cancel ongoing harm.





Safety Comes Before Repairing the Relationship

If a relationship includes hitting, threats, stalking, coercion, financial control, or isolation from friends and family, the priority is no longer better communication or healing the other person. The priority is your safety. When possible, contact trusted people and qualified local services, preserve necessary evidence, and make a safety plan under conditions that do not increase danger. Avoid confronting the person alone if doing so could escalate.

Being moved by vulnerability may come from empathy, familiarity, or a desire to feel needed. That does not make your feelings false, and it does not mean every difficult relationship must end immediately. It does invite a clearer question: Am I loving a real person who is willing to take responsibility, or am I protecting the hope that everything will change if I work hard enough?

Healthy intimacy is not one person continually burning so the other can wait to be illuminated. A more stable relationship allows both people to have ground of their own and to nurture each other. You can care while keeping boundaries. You can give and receive. You can understand someone's past without becoming fully responsible for their present.



中意人寿
GENERALI CHINA

GCL Customer Service Hotline: 400 888 7555